

A NONLINEAR EVOLUTION EQUATION WITH 2 + 1 DIMENSIONS RELATED TO NONSTATIONARY DIRAC-TYPE SYSTEM

MANSUR I ISMAILOV

ABSTRACT. In this paper the inverse scattering problem for the nonstationary Dirac-type system on the whole plane was considered. A nonlinear evolution sytem of equation related to nonstationary Dirac-type system is introduced and the solvibility of this sytem using the IST method is studied.

1. INTRODUCTION

As is known, every linear scattering problem determines a class of nonlinear evolution equations for which the inverse scattering transform (IST) method is suitable to integrate these equations. The investigation of the nonlinear evolution equations by the IST method has been stated in [1, 2]. The essence of this method is to represent this nonlinear equation as a Lax equation

$$(0.1) \quad \frac{\partial \mathbf{L}}{\partial t} = [\mathbf{L}, \mathbf{A}]$$

by using the Lax pair $\mathbf{L}$ and $\mathbf{A}$. The equation in the form of (0.1) allows to investigate evolution of scattering data instead of the evolution of the operator $\mathbf{L}$, if the inverse scattering problem is investigated for the operator $\mathbf{L}$.

The IST method for the nonlinear evolution equations with 1+1 dimensions (one space and one time dimensions) have been described in a variety reviews and monographs. The generalization of the IST method to nonlinear evolution equations with 2+1 dimensions (two space and one time dimension) has been developed in monograph [3]. In this direction one can also notice the papers [4 - 7], in which the inverse scattering problem for the first order systems is studied and this inverse problem was applied to integration of N - wave interactions, Davey-Stewartson and Kodomtsev-Petviashvili equations.

Let $\mathbf{L} = \frac{\partial}{\partial y} - \mathbf{M}$, where $\mathbf{M}$ is an ordinary differential operator in x . Then equation (0.1) is rewritten as (see [2])

$$(0.2) \quad \left[\frac{\partial}{\partial y} - \mathbf{M}, \frac{\partial}{\partial t} - \mathbf{A} \right] = 0$$

Date: 15, August, 2009.

2000 Mathematics Subject Classification. Primary 35R30 ; Secondary 35L50, 35P25, 37K15.

Key words and phrases. Nonstationary scattering problem, Dirac-type system, Inverse problem on the plane, Nonlinear evolution equations.

and it is a relation for ordinary differential operators. Some nonlinear evolution equations in 2+1 dimension are presented in [8] by using commutativity condition (0.2) when $\mathbf{M}$ is an scalar coefficients ordinary differential operator in x .

This article is organized as follows: In Section 2, we consider the inverse scattering problems for the $\mathbf{L} = \frac{\partial}{\partial y} - \mathbf{M}$, with $\mathbf{M} = \sigma \frac{\partial}{\partial x} + Q(x, y)$ in the whole plane. We introduce the scattering data for the operator $\mathbf{L}$ in the plane (the minimal information for a unique restoration of matrix-function $Q(x, y)$). All the results in Section 2 are the matrix generalization of the L. P. Nizhnik's results for two component nonstationary Dirac system on the whole plane in [9] (see [4] also). Taking into account that this generalization is not difficult, so we do not give the proofs. In Section 3, we introduce some system of nonlinear evolution equations with 2+1 dimensions related to nonstationary Dirac-type systems and we study the solvability of this system using the IST method. In [10], the IST method is applied to integrate the two component nonlinear Schrödinger equation, using the inverse scattering problem for stationary Dirac-type system in whole line.

2. Inverse scattering problem for the Dirac-type system in the plane

Let us consider a system of first-order partial differential equations (PDEs)

$$(1.1) \quad \mathbf{L}(\psi) \equiv \frac{\partial \psi}{\partial y} - \sigma \frac{\partial \psi}{\partial x} - Q(x, y) \psi = 0, \quad -\infty < x, y < +\infty,$$

where

$$\sigma = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad Q = \begin{bmatrix} 0 & 0 & q_1 \\ 0 & 0 & q_2 \\ q_3 & q_4 & 0 \end{bmatrix}.$$

with the measurable complex-valued functions q_i ($i = 1, \dots, 4$).

We give fundamental results on the inverse scattering problem for the system (1.1) in the case where the coefficients q_i ($i = 1, \dots, 4$) decrease quite fast with respect to variables x and y at infinity. Each bounded solution of the system (1.1) has the asymptotic

$$\begin{aligned} \psi(x, y) &= F_x a_-(y) + o(1), \quad y \longrightarrow -\infty, \\ \psi(x, y) &= F_x a_+(y) + o(1), \quad y \longrightarrow +\infty. \end{aligned}$$

where F_x denotes the diagonal matrix shift operator, such that for a vector function $a(y) = \text{col}[a_1(y), a_2(y), a_3(y)]$, $F_x a(y) = \text{col}[a_1(t+x), a_2(t+x), a_3(t-x)]$

The scattering operator S is defined by the equality

$$(1.2) \quad a_+(y) = \mathbf{S} a_-(y).$$

This operator has the inverse S^{-1} and $S = I + F$ and $S^{-1} = I + G$, where $F = [F_{ij}]_{i,j=1}^3$ and $G = [G_{ij}]_{i,j=1}^3$ are matrix integral operators. To regenerate the system (1.1), i.e., the coefficients q_i ($i = 1, \dots, 4$) it is sufficient to know F_{13}, F_{23}, G_{31} and G_{32} . The collection $\{F_{13}, F_{23}, G_{31}, G_{32}\}$ is called the scattering data for the system (1.1) in the plane. The solution of the inverse problem reduces to the integral equation

$$A(x, y, \tau) - \int_{-\infty}^y \left[\int_y^{+\infty} A(x, y, z) \tilde{G}(z - x, s + x) dz \right] \tilde{F}(s + x, \tau - x) ds = \tilde{F}(y + x, \tau - x), \quad (1.3)$$

$$B(x, y, \tau) - \int_y^{+\infty} \left[\int_{-\infty}^y B(x, y, z) \tilde{F}(z + x, s - x) dz \right] \tilde{G}(s + x, \tau - x) ds = \tilde{G}(y - x, \tau + x),$$

where $\tilde{F} = \begin{bmatrix} F_{13} \\ F_{23} \end{bmatrix}$ and $\tilde{G} = \begin{bmatrix} G_{31} & G_{32} \end{bmatrix}$.

The coefficients q_i ($i = 1, \dots, 4$) of the system (1.1) are expressed in terms of the solution of (1.3) by means of the equations

$$\begin{bmatrix} q_1 \\ q_2 \end{bmatrix} (x, y) = -2A(x, y, y), \quad \begin{bmatrix} q_3 & q_4 \end{bmatrix} (x, y) = -2B(x, y, y).$$

3. NONLINEAR EVOLUTION EQUATION RELATED TO NONSTATIONARY DIRAC-TYPE SYSTEMS

Let $\mathbf{M}$ and $\mathbf{A}$ be first order matrix operators in (0.2):

$$\mathbf{M} = \sigma \frac{\partial}{\partial x} + Q, \quad \mathbf{A} = \tau \frac{\partial}{\partial x} + P.$$

Then the equation (0.2) becomes to the form

$$(2.1) \quad \left[\frac{\partial}{\partial y} - \sigma \frac{\partial}{\partial x} - Q, \frac{\partial}{\partial t} - \tau \frac{\partial}{\partial x} - P \right] = 0.$$

Here τ and P are third order square matrices. Let the matrix τ be real and diagonal: $\tau = \text{diag}(b_1, b_2, b_3)$, $b_1 > b_2 > b_3$ and the matrices Q and P obey the relation: $[\sigma, P] = [\tau, Q]$.

Take $P = \begin{bmatrix} 0 & v_{12} & \frac{b_1 - b_3}{2} q_1 \\ v_{21} & 0 & \frac{b_2 - b_3}{2} q_2 \\ \frac{b_1 - b_3}{2} q_3 & \frac{b_2 - b_3}{2} q_4 & 0 \end{bmatrix}$, where the functions v_{12} and v_{21} are solutions of the equations

$$\begin{aligned} \frac{\partial}{\partial y} v_{12} - \frac{\partial}{\partial x} v_{12} &= -\frac{b_1 - b_2}{2} q_1 q_4, \\ \frac{\partial}{\partial y} v_{21} - \frac{\partial}{\partial x} v_{21} &= -\frac{b_2 - b_1}{2} q_2 q_3, \end{aligned}$$

respectively.

Then, the conditions (2.1) are reduced to the system of equations

$$\begin{aligned} \frac{\partial}{\partial t} q_1 + k_1 \frac{\partial}{\partial y} q_1 + k_2 \frac{\partial}{\partial x} q_1 &= v_{12} q_2, \\ \frac{\partial}{\partial t} q_2 + k_1 \frac{\partial}{\partial y} q_2 + k_2 \frac{\partial}{\partial x} q_2 &= v_{21} q_1, \\ \frac{\partial}{\partial t} q_3 + k_3 \frac{\partial}{\partial y} q_3 + k_4 \frac{\partial}{\partial x} q_3 &= -v_{21} q_4, \\ \frac{\partial}{\partial t} q_4 + k_3 \frac{\partial}{\partial y} q_4 + k_4 \frac{\partial}{\partial x} q_4 &= -v_{12} q_3, \end{aligned} \quad (2.2)$$

where $k_1 = -\frac{b_1-b_3}{2}$, $k_2 = -\frac{b_1+b_3}{2}$, $k_3 = -\frac{b_2-b_3}{2}$, $k_4 = -\frac{b_2+b_3}{2}$.

Let us denote $\mathbf{P} = \frac{\partial}{\partial t} - \tau \frac{\partial}{\partial y} - P$. The Lax form (2.1) of the system of equations (2.2) enables us to apply to IST method for the integration.

Lemma 1. *Let ψ be a solution of the Dirac-type system (1.1), whose the coefficients $q_i, i = 1, \dots, 4$ satisfy Eqs. (2.2). Then the function $\varphi = \mathbf{P}\psi$ also satisfy the system (1.1).*

Proof. Let us apply the operator equation to ψ :

$$(\mathbf{L}\mathbf{P} - \mathbf{P}\mathbf{L})\psi = \mathbf{L}(\mathbf{P}\psi) - \mathbf{P}(\mathbf{L}\psi) = \mathbf{0}.$$

Taking into account that $\mathbf{L}\psi = \mathbf{0}$, then

$$\mathbf{L}(\mathbf{P}\psi) = \mathbf{0}$$

follows from the last equation. It means that $\mathbf{P}\psi$ is solution of system (1.1). $\square$

The next theorem is true for the evolution of the scattering data.

Lemma 2. *Let the coefficients $q_i, i = 1, \dots, 4$ of the Dirac-type system (1.1) depend on t as a parameter and satisfy the system of equation (2.2). Besides that*

$$(2.3) \quad v_{12}(x, \pm\infty) = 0, \quad v_{21}(x, \pm\infty) = 0.$$

Then the kernels $F_{13}(y, \tau; t), F_{23}(y, \tau; t), G_{31}(y, \tau; t), G_{32}(y, \tau; t)$ of the integral operators $F_{13}, F_{23}, G_{31}, G_{32}$ corresponding to the scattering operator $\mathbf{S}$ for the system (1.1) in the plane satisfy the system of equations

$$\begin{aligned} \frac{\partial}{\partial t} F_{13}(y, \tau; t) - b_1 \frac{\partial}{\partial y} F_{13}(y, \tau; t) + b_3 \frac{\partial}{\partial \tau} F_{13}(y, \tau; t) &= 0, \\ \frac{\partial}{\partial t} F_{23}(y, \tau; t) - b_2 \frac{\partial}{\partial y} F_{23}(y, \tau; t) + b_3 \frac{\partial}{\partial \tau} F_{23}(y, \tau; t) &= 0 \end{aligned} \quad (2.4)$$

and

$$\begin{aligned} \frac{\partial}{\partial t} G_{31}(y, \tau; t) + b_3 \frac{\partial}{\partial y} G_{31}(y, \tau; t) - b_1 \frac{\partial}{\partial \tau} G_{31}(y, \tau; t) &= 0, \\ \frac{\partial}{\partial t} G_{32}(y, \tau; t) + b_3 \frac{\partial}{\partial y} G_{32}(y, \tau; t) - b_2 \frac{\partial}{\partial \tau} G_{32}(y, \tau; t) &= 0. \end{aligned} \quad (2.5)$$

Proof. By virtue of definition of the scattering operator $\mathbf{S}$, from Lemma 1 we get

$$(2.6) \quad \varphi_+ = \mathbf{S}\varphi_-$$

where $\varphi_{\pm} = \mathbf{P}a_{\pm}$, $\mathbf{P} = \frac{\partial}{\partial t} - \tau \sigma \frac{\partial}{\partial y}$.

Since $a_+ = \mathbf{S}a_-$ (see (1.2)), the equality

$$(2.7) \quad \mathbf{P}\mathbf{S} = \mathbf{S}\mathbf{P}$$

follows from (2.6).

Analogously,

$$(2.8) \quad \mathbf{P}\mathbf{S}^{-1} = \mathbf{S}^{-1}\mathbf{P}.$$

From the matrix operator equation (2.7) it follows that the kernels of the integral operators F_{13} and F_{23} satisfy system of equation (2.4). The similarly system of

equation (2.5) for the kernels of the integral operators G_{31} and G_{32} follows from the matrix operator equation (2.8). $\square$

According to Chapter 2 and Lemma 2, let us give a procedure for the solution of the system (2.2) by IST method.

Theorem 1. *The system of equations (2.2) admits integration in the class of decreasing functions that satisfy the condition (2.3) by the IST method. The solution $q_i, i = 1, \dots, 4$ of the system (2.2) is determined by formulae*

$$\begin{bmatrix} q_1 \\ q_2 \end{bmatrix} (x, y; t) = -2A(x, y, y; t), \quad \begin{bmatrix} q_3 & q_4 \end{bmatrix} (x, y; t) = -2B(x, y, y; t),$$

where the vector functions A and B are the solutions of the integral equations

$$\begin{aligned} A(x, y, \tau; t) - \int_{-\infty}^y \left[\int_y^{+\infty} A(x, y, z; t) \tilde{G}(z - x, s + x; t) dz \right] \tilde{F}(s + x, \tau - x; t) ds \\ = \tilde{F}(y + x, \tau - x; t), \\ B(x, y, \tau; t) - \int_y^{+\infty} \left[\int_{-\infty}^y B(x, y, z; t) \tilde{F}(z + x, s - x; t) dz \right] \tilde{G}(s + x, \tau - x; t) ds \\ = \tilde{G}(y - x, \tau + x; t), \end{aligned}$$

where $\tilde{F} = \begin{bmatrix} F_{13} \\ F_{23} \end{bmatrix}$ and $\tilde{G} = \begin{bmatrix} G_{31} & G_{32} \end{bmatrix}$ are satisfied the evolution equations (2.4) and (2.5).

Thus the system of nonlinear evolution equation (2.2) admits the integration by IST method. Let us denote by Π the operator transforming the coefficients $q(t) = \text{col}(q_1(x, y; t), q_2(x, y; t), q_3(x, y; t), q_4(x, y; t))$ of the equation (1.1) to scattering data $T(t) = \text{col}(F_{13}(y, \tau; t), F_{23}(y, \tau; t), G_{31}(y, \tau; t), G_{32}(y, \tau; t))$ as follows:

$$\Pi : q(t) \mapsto T(t).$$

Then the solution of the system (2.2) can be represent as

$$(2.9) \quad q(t) = \Pi^{-1} e^{t\mathbf{M}} T$$

where $T = T(0)$ i.e. $T = \text{col}(F_{13}(y, \tau), F_{23}(y, \tau), G_{31}(y, \tau), G_{32}(y, \tau))$ and

$$\mathbf{M} = \begin{bmatrix} b_1 \frac{\partial}{\partial y} - b_3 \frac{\partial}{\partial \tau} & 0 & 0 & 0 \\ 0 & b_2 \frac{\partial}{\partial y} - b_3 \frac{\partial}{\partial \tau} & 0 & 0 \\ 0 & 0 & b_1 \frac{\partial}{\partial \tau} - b_3 \frac{\partial}{\partial y} & 0 \\ 0 & 0 & 0 & b_2 \frac{\partial}{\partial \tau} - b_3 \frac{\partial}{\partial y} \end{bmatrix}.$$

The formula (2.9) is a specified form of solution $q(t)$ of (2.2).

REFERENCES

- [1] M. J. Ablowitz and H. Segur, *Solitons and the inverse scattering transform*. SIAM Stud. Appl. Math. **4**, Philadelphia, 1981.
- [2] V. E. Zakharov, S. V. Manakov, S. P. Novikov, and L. P. Pitaevski, *Theory of solitons. The inverse scattering method*. Nauka, Moscow, 1980 (in Russian); English transl. Consultants bureau (New York: Plenum) 1984.
- [3] B. G. Konopelchenko, *Introduction to multidimensional integrable equations. The inverse spectral transform in 2+1 dimensions*. Plenum Press, New York, 1992. 292 pp.
- [4] L. P. Nizhnik, *The inverse scattering problem for hyperbolic equations and their application to nonlinear integrable systems*. Reports on Math. Phys. **26** (2), 261-283 (1988).

- [5] L. P. Nizhnik and M. D. Pachinaiko, *Integration of the nonlinear two-dimensional spatial Schrödinger equation by the inverse-problem method*, Functional Anal. Appl., 1982, **16** (1), 66-69
- [6] A. S. Fokas and L. Y. Sung, *On the solvability of the N- Wave, Davey-Stewartson and Kodomtsev-Petviashvili equations*, Inverse Problems, 1992, **8**, 673-708.
- [7] A. S. Fokas and M. J. Ablowitz, *On the inverse scattering transform of multidimensional nonlinear equations related to first-order systems in the plane*, J. Math. Phys., 1984, **25** (8), 2494-2505.
- [8] B. G. Konopelchenko and V. G. Dubrovsky, *Some new integrable nonlinear evolution equations in 2+1 dimensions*. Phys. Lett. A 102 (1984), no. 1-2, 15-17
- [9] L. P. Nizhnik, *Inverse nonstationary scattering problems*. Nauk. Dumka, Kiev, 1973 (in Russian).
- [10] S. V. Manakov, *On the theory of two-dimensional stationary self-focusing of electromagnetic waves*, Sov. Phys. JETP 38, 248-253 (1974).

DEPARTMENT OF MATHEMATICS, GEBZE INSTITUTE OF TECHNOLOGY, GEBZE-KOCAELI 41400, TURKEY

E-mail address: `mismailov@gyte.edu.tr`